

Review Article

Avascular Blebs and Late Bleb Leakage; A Review of Causes and Management

Maryam Yadgari ^{1,2}, MD; Kiana Hassanpour ^{*1,2} MD, MPH

1. Imam Hossein Medical Center, Ophthalmology Department, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Ophthalmology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

***Corresponding Author:** Kiana Hassanpour

E-mail: kiana.hassanpour@gmail.com

Article Notes:

Received: May. 9, 2018

Received in revised form:
Jul. 2, 2018

Accepted: Aug. 18, 2018

Available Online: Sep. 29,
2018

Keywords:

Trabeculectomy

Mitomycin

Leakage

Avascular

Abstract

Trabeculectomy with mitomycin-C remains the gold standard for surgical glaucoma management; however, this technique includes some sight-threatening complications like avascular thin bleb, subsequent leakage and ultimately endophthalmitis. To date, various non surgical methods have been reported for the management of bleb leakage, but surgical management frequently becomes necessary especially in frank leakages. The most common surgical approach includes excision of the avascular and necrotic leaking bleb combined with the advancement of adjacent healthy conjunctiva. The aim of the present review is to discuss avascular bleb and late bleb leakage after trabeculectomy including their histopathology, risk factors, prevention and management.

How to cite this article: Yadgari M, Hassanpour K. Avascular Blebs and Late Bleb Leakage; A Review of Causes and Management . Journal of Ophthalmic and Optometric Sciences . 2018;2(4): 31-9.

Introduction

The ultimate goal of trabeculectomy is the prevention of glaucomatous optic neuropathy progression through intraocular pressure (IOP) reduction and without development of hypotony¹. This result is achieved via a diffuse filtering bleb formation best made by a fornix-based conjunctival flap rather than limbal-based techniques².

Mitomycin C (MMC), an antimetabolite agent isolated from *Streptomyces Caespitosus*, prevents DNA synthesis and reduces the chance of fibrosis³. Although it has been widely used during trabeculectomy to prevent fibrosis and increase surgical success, it serves as a double sword that can produce thin and avascular blebs with ischemia and leakage as the possible inadvertent consequences⁴⁻¹⁰.

A large avascular area of the bleb is the most common predisposing factor for late bleb leakage compared with a cystic bleb¹¹. Moreover, most avascular blebs show avascularity within the first postoperative year with the anterior location to be the most common site¹². An avascular bleb could result in the late-onset bleb leakage months to years after trabeculectomy, which is directly proportional to the duration of avascularity^{11,13-19}. Bleb leakage increases the risk of bleb-associated blebitis and endophthalmitis up to 26 folds, as the most catastrophic event after trabeculectomy²⁰. In a study by Wolner et al.,²¹ 62 percent of the patients with bleb-related endophthalmitis had chronic and intermittent leakage. Interestingly all these patients had thin-walled blebs²¹. The incidence of late-onset bleb leakage after trabeculectomy with adjunctive antimetabolites varies between 1.8 % and 12.9 %^{10, 22-24}.

Conjunctival epithelial defect mainly predisposes the underlying bleb to frank late-onset bleb leaks²⁵. The possible mechanism

is a higher chance of aqueous oozing through the unhealed epithelial defect that can eventually deteriorate into an obvious point leakage²⁵. Moreover, avascular blebs can cause potentially sight-threatening hypotony-related complications including flat anterior chamber, peripheral anterior synechiae, cataract, choroidal detachment and hypotonic maculopathy^{10, 26-31}.

The aim of the present review is to discuss avascular bleb and late bleb leakage after trabeculectomy including their histopathology, risk factors, prevention and management.

Methods

A PubMed and Google scholar search was performed, analyzing all publications from 1970 to 2018 regarding the topic “avascular bleb and late-onset bleb leakage” using keywords: bleb, late onset, leakage, avascular, and hypovascular. Only studies published in English language were included for this review.

Histopathology

Studies investigating histopathological characteristics of avascular blebs show the presence of irregular conjunctival epithelium, multiple breaks in the basement membrane, irregularly arranged collagen with few fibroblasts and lack of vessels in the substantia propria^{6, 32-34}. Furthermore, the electron microscopy study of avascular blebs shows microvilli at the superficial area in some parts of the conjunctival epithelium⁶. Late-leaking blebs also contain focal epithelial thinning and interruption, hypocellularity, and stromal collagen degeneration¹⁰.

Risk factors

The prevalence of bleb leakage among patients undergoing trabeculectomy with antimetabolites is estimated to be from 1.3 %

to 25 %^{14, 35-38}. Avascular bleb formation is more frequent in a limbus-based conjunctival flap compared with a fornix-based flap^{2, 39}. Excessive and sustained aqueous filtration is another risk factor for this complication^{34, 40}. Conversely, a diffuse vascularized bleb with wide posterior extension decreases the risk of bleb failure and leakage²⁴. Another mechanism causing bleb avascularity is the formation of a localized bleb with subsequent increase in the internal pressure of the bleb³⁴.

Prevention

There is no exact method to prevent the bleb avascularity⁴⁰. Replacing MMC with another agent to lower complications has been tried in various studies. Nilforoushan et al.,⁴¹ compared Bevacizumab with MMC in trabeculectomy and showed similar complication rates between the two drugs and higher postoperative glaucoma medications needed in the Bevacizumab group. In two other studies, the same author investigated combined use of trabeculectomy and OculusGen implanted under the bleb and reported that this method was safe but still required higher glaucoma medications compared to MMC-trabeculectomy^{42, 43}.

Okuda et al.,⁴⁰ found that application of a biodegradable, honeycomb-patterned film reduces bleb avascularity without affecting filtration surgery success rate in rabbits undergoing trabeculectomy with MMC by avoiding conjunctival exposure to excess filtrated aqueous⁴⁰.

Management

A thin avascular bleb can be closely followed. Various alarm signs including the presence of a shallow anterior chamber, reduced visual acuity, development of hypotony maculopathy, choroidal detachment and more importantly

frank bleb leakage necessitate treatment^{12, 18}. Bleb leakage needs special attention because of increased risk of endophthalmitis besides other hypotony related complications^{13, 14, 21, 38, 44-46}. The ultimate goal of bleb leakage treatment includes the elimination of leakage as well as preserving filtration bleb function⁴⁷.

Conservative Management

Multiple modalities have been used for conservative management of bleb leakage including sodium hyaluronate and autologous serum eye drop²⁵, vigorous treatment of concomitant dry eye^{1,48}, suppressants of aqueous humor production⁴⁹, bandage contact lenses^{49,50}, tissue adhesives⁵⁰⁻⁵⁴, induction of an inflammatory healing response in the bleb with chemical irritants⁵³, and symblepharon rings⁵⁵.

Surgical Management

Several techniques for surgical management of a leaking bleb have been reported with most of them reserved as the last resort. However, the surgical approach becomes necessary in many patients. These approaches include covering the leakage area with an overlying conjunctival flap⁵⁶⁻⁵⁸, external conjunctival cryopexy⁵⁹, thermal coagulation by cautery⁶⁰, application of argon laser⁶¹, collagen crosslinking (CXL)^{62,63}, direct suturing of the conjunctival defect (so-called buttonhole)^{64,65}, patching the sclerostomy site with a corneal⁶⁶ or scleral tissue^{41,67-69}, advancing a flap of Tenon's capsule from the superior fornix to plug the leakage area^{56, 57, 70}, surgical revision with the advancement of fornix conjunctival flap^{5, 46, 65, 71, 72}, free conjunctival autograft from either the same or the fellow eye⁷³, compression sutures⁷⁴, intra bleb autologous blood injection^{75, 76}, reconstruction with preserved human amniotic membrane for the

repair of difficult bleb leaks^{51, 77}, and bleb needling techniques⁷⁸.

Direct manipulation of a thin avascular bleb can cause traumatic damages to a semi-destroyed area that can worsen the bleb leakage. Thereby bleb revision surgeries are often considered as the last option to avoid bleb failure⁷⁹.

Regarding chronic leakages, there is often an epithelial tract that originates from bleb's surface entering into the episclera. To eliminate the leakage, this tract must be completely excised³¹. Since the leaking area is usually surrounded by a thin tissue, most of these procedures might be complicated by more damage to the bleb⁴⁹.

Currently, the most recommended surgical technique is excision of the leaking area of the bleb and the advancement of a fornix-based conjunctival flap made through undermining the surrounding conjunctiva combined with the widespread use of relaxing incisions to avoid a tight flap^{46, 65, 71, 72}. In this technique, the anterior margin of the flap is sutured into the cornea. Nevertheless, 44 % to 86 % of patients undergoing bleb repair will require glaucoma medications for IOP control⁷⁹.

Discussion

Intraoperative antifibrotic agents like MMC are routinely used in trabeculectomy²⁸. The main drawback of their usage is the increased risk of avascularity and thinning of the blebs. The risk of late bleb leakage is 3 times higher with the application of MMC compared with 5-fluorouracil¹. Some blebs will become progressively thin, oversized, and vulnerable; therefore they are at risk of recurrent leakages and over filtration^{6, 10}. Moreover, the 5-year risk of bleb related endophthalmitis, the most terrible complication of trabeculectomy, is increased up to 23 % after bleb leakage^{9, 80}. Conjunctival epithelial defect predisposes bleb

to frank late-onset bleb leaks through chronic aqueous oozing³¹. Therefore, maintaining an effective tear film is crucial for inhibiting a possible progression of epithelial defect into a dangerous bleb leakage²⁵. Ophthalmologists should distinguish frank leakage containing an obvious site from bleb sweating as areas of pinpoint transconjunctival flow since the latter usually is not correlated with severe complications⁶.

Many medically treated leaking blebs finally recur after conservative managements like the use of sodium hyaluronate and autologous serum^{6, 25, 31}. In longstanding leakages, the development of an epithelial tract in the leaking area would complicate the medical management and necessitates the surgical approach^{21, 25}. The use of prophylactic topical antibiotics to minimize the risk of endophthalmitis is a controversial issue because of the increased risk of bacterial flora resistance that could result in a more aggressive infection²⁴.

Excision of the leaking area followed by the advancement of a conjunctival flap is the most recommended surgical approach^{46, 65, 71, 72}. To prevent the development of a tight or small flap, some surgeons replace the excised necrotic area with a free conjunctival graft borrowed from ipsilateral side or the fellow eye⁴⁷. These surgical modalities all share the risk of bleb failure⁴⁹. van de Geijn et al.,⁷⁹ in a retrospective study of 36 cases, reported three failed filtering blebs within 1 month and 2 more cases within 1 year after surgery because of subconjunctival and episcleral scarring. Complete and qualified success rates in their study were 51.6 % and 86.1 %, respectively and failed cases finally underwent re-trabeculectomy⁷⁹. One eye was complicated with conjunctival and Tenon flap dehiscence and one eye was affected by

conjunctival wound leak at the limbus which underwent suturing⁷⁹. Avascular leaking bleb recurred in one eye 19 months after surgery⁷⁹. In a retrospective study Al-Shahwan et al.,⁸¹ studied 34 eyes that underwent bleb excision with conjunctival advancement for late bleb leakage after trabeculectomy and achieved IOP control without medication in 58.8 % of the operated eyes. Early leakage, late recurrent or persistent leakage, and endophthalmitis occurred in 7, 1 and 1 cases, respectively⁸¹. Lee et al.,⁸² in a retrospective review of 17 eyes undergoing avascular bleb excision with conjunctival advancement reported a success rate of 100 % with IOP between 6 and 18 mmHg with or without medication at 51 months of follow up. They reported relapse with perforation of the avascular area in one case 89 months after surgery⁸². Blepharoptosis occurred in 23 % of their cases in whom spontaneous resolution was finally achieved and only one case with sustained hypertropia occurred⁸². Budenz et al.,⁸³ compared amniotic membrane transplantation with conjunctival advancement for the repair of late-onset bleb leakage and reported seven failures in the amniotic membrane transplant group versus no failure in the conjunctival advancement group. Of those seven patients with failures, two had persistent leaks that were unresponsive to further suturing, two had late-onset leakages,

and three required repeated glaucoma surgery⁸³. The cumulative survival for amniotic membrane transplant was 81 % at six months, 74 % at one year, and 46 % at two years, compared with 100 % in conjunctival advancement group⁸³. Complications of bleb excision with conjunctival advancement include postoperative IOP spike, bleb leakage recurrence, bleb dysesthesia, filtering bleb failure, blepharoptosis, and hypertropia⁸⁴⁻⁸⁷. Nevertheless, this procedure seems to be the best surgical treatment for complicated avascular blebs that need an operation⁴⁷.

Conclusion

Avascular bleb and its potential complications should be assessed at each visit after trabeculectomy. Early detection may increase the chance of success for conservative treatments. Ophthalmologists should rigorously follow cases that are at risk for bleb associated infections including patients with frank leakage and should use surgical modalities to prevent potentially catastrophic sight-threatening events.

Authors ORCIDs

Maryam Yadgari:

 <https://orcid.org/0000-0003-0829-1861>

Kiana Hassanpour:

 <https://orcid.org/0000-0002-1788-7352>

References

1. Liang SYW, Lee GA, Whitehead K. Histopathology of a functioning mitomycin-C trabeculectomy. *Clinical & experimental ophthalmology*. 2009;37(3):316-9.
2. Agbeja AM, Dutton GN. Conjunctival incisions for trabeculectomy and their relationship to the type of bleb formation--a preliminary study. *Eye (Lond)*. 1987;1 (Pt 6):738-43.
3. Al Habash A, Aljasim LA, Owaidhah O, Edward DP. A review of the efficacy of mitomycin C in glaucoma filtration surgery. *Clin Ophthalmol*. 2015;9:1945-51.
4. Greenfield DS, Liebmann JM, Jee J, Ritch R. Late-onset bleb leaks after glaucoma filtering surgery. *Arch Ophthalmol*. 1998;116(4):443-7.

5. Shields MB, Scroggs MW, Sloop CM, Simmons RB. Clinical and histopathologic observations concerning hypotony after trabeculectomy with adjunctive mitomycin C. *Am J Ophthalmol.* 1993;116(6):673-83.
6. Nuyts RM, Felten PC, Pels E, Langerhorst CT, Geijssen HC, Grossniklaus HE, et al. Histopathologic effects of mitomycin C after trabeculectomy in human glaucomatous eyes with persistent hypotony. *Am J Ophthalmol.* 1994;118(2):225-7.
7. Greenfield DS, Parrish RK 2nd. Bleb rupture following filtering surgery with mitomycin-C: clinicopathologic correlations. *Ophthalmic Surg Lasers.* 1996;27(10):876-7.
8. Sihota R, Dada T, Gupta SD, Sharma S, Arora R, Agarwal HC. Conjunctival dysfunction and mitomycin C-induced hypotony. *J Glaucoma.* 2000;9(5):392-7.
9. DeBry PW, Perkins TW, Heatley G, Kaufman P, Brumback LC. Incidence of late-onset bleb-related complications following trabeculectomy with mitomycin. *Arch Ophthalmol.* 2002;120(3):297-300.
10. Belyea DA, Dan JA, Stamper RL, Lieberman MF, Spencer WH. Late onset of sequential multifocal bleb leaks after glaucoma filtration surgery with 5-fluorouracil and mitomycin C. *Am J Ophthalmol.* 1997;124(1):40-5.
11. Hu C-Y, Matsuo H, Tomita G, Suzuki Y, Araie M, Shirato S, et al. Clinical characteristics and leakage of functioning blebs after trabeculectomy with mitomycin-C in primary glaucoma patients. *Ophthalmology.* 2003;110(2):345-52.
12. Kohl DA, Walton DS. Limbus-based versus fornix-based conjunctival flaps in trabeculectomy: 2005 update. *Int Ophthalmol Clin.* 2005;45(4):107-13.
13. Khaw PT, Doyle JW, Sherwood MB, Smith MF, McGorray S. Effects of intraoperative 5-fluorouracil or mitomycin C on glaucoma filtration surgery in the rabbit. *Ophthalmology.* 1993;100(3):367-72.
14. Ticho U, Ophir A. Late complications after glaucoma filtering surgery with adjunctive 5-fluorouracil. *Am J Ophthalmol.* 1993;115(4):506-10.
15. Gressel MG, Parrish II RK, Folberg R. 5-fluorouracil and glaucoma filtering surgery: I. An animal model. *Ophthalmology.* 1984;91(4):378-83.
16. Schwartz AL, Weiss HS. Bleb leak with hypotony after laser suture lysis and trabeculectomy with mitomycin C. *Arch Ophthalmol.* 1992;110(8):1049.
17. Blondeau P, Phelps CD. Trabeculectomy vs thermosclerostomy. A randomized prospective clinical trial. *Arch Ophthalmol.* 1981;99(5):810-16.
18. Lamping KA, Bellows AR, Hutchinson BT, Afran SI. Long-term evaluation of initial filtration surgery. *Ophthalmology.* 1986;93(1):91-101.
19. Anand N, Arora S, Clowes M. Mitomycin C augmented glaucoma surgery: evolution of filtering bleb avascularity, transconjunctival oozing, and leaks. *Br J Ophthalmol.* 2006;90(2):175-80.
20. Soltau JB, Rothman RF, Budenz DL, et al. Risk factors for glaucoma filtering bleb infections. *Arch Ophthalmol.* 2000;118(3):338-42.
21. Wolner B, Liebmann JM, Sassani JW, Ritch R, Speaker M, Marmor M. Late bleb-related endophthalmitis after trabeculectomy with adjunctive 5-fluorouracil. *Ophthalmology.* 1991;98(7):1053-60.
22. Matsuo H, Tomidokoro A, Suzuki Y, Shirato S, Araie M. Late-onset transconjunctival oozing and point leak of aqueous humor from filtering bleb after trabeculectomy. *Am J Ophthalmol.* 2002;133(4):456-62.
23. Caronia RM, Liebmann JM, Friedman R, Cohen H, Ritch R. Trabeculectomy at the inferior limbus. *Arch Ophthalmol.* 1996;114(4):387-91.
24. Alwitary A, King AJ. Surveillance of late-onset bleb leak, blebitis and bleb-related endophthalmitis—a UK incidence study. *Graefes Arch Clin Exp Ophthalmol.* 2012;250(8):1231-6.
25. Sagara H, Iida T, Saito K, Noji H, Ogasawara M, Oyamada H. Conservative treatment for late-onset bleb leaks after trabeculectomy with mitomycin C in patients with ocular surface disease. *Clin Ophthalmol.* 2012;6:1273-9.
26. Mandelbaum S, Forster RK, Gelender H, Culbertson W. Late onset endophthalmitis associated with filtering blebs. *Ophthalmology.* 1985;92(7):964-



- 72.
27. Waheed S, Liebmann JM, Greenfield DS, et al. Recurrent bleb infections. *Br J Ophthalmol*. 1998;82(8):926-9.
28. Mochizuki K, Jikihara S, Ando Y, Hori N, Yamamoto T, Kitazawa Y. Incidence of delayed onset infection after trabeculectomy with adjunctive mitomycin C or 5-fluorouracil treatment. *Br J Ophthalmol*. 1997;81(10):877-83.
29. Hattenhauer JM, Lipsich MP. Late endophthalmitis after filtering surgery. *Am J Ophthalmol*. 1971;72(6):1097-101.
30. Poulsen EJ, Allingham RR. Characteristics and risk factors of infections after glaucoma filtering surgery. *J Glaucoma*. 2000;9(6):438-43.
31. Sagara H, Iida T, Suzuki K, Fujiwara T, Koizumi H, Yago K. Sodium hyaluronate eye drops prevent late-onset bleb leakage after trabeculectomy with mitomycin C. *Eye (Lond)*. 2008;22(4):507-14.
32. Mietz H, Arnold G, Kirchhof B, Diestelhorst M, Krieglstein GK. Histopathology of episcleral fibrosis after trabeculectomy with and without mitomycin C. *Graefes Arch Clin Exp Ophthalmol*. 1996;234(6):364-8.
33. Hutchinson AK, Grossniklaus HE, Brown RH, McManus PE, Bradley CK. Clinicopathologic features of excised mitomycin filtering blebs. *Arch Ophthalmol*. 1994;112(1):74-9.
34. Hirooka K, Mizote M, Baba T, Takagishi M, Shiraga F. Risk factors for developing avascular filtering bleb after fornix-based trabeculectomy with mitomycin C. *J Glaucoma*. 2009;18(4):301-4.
35. Rockwood EJ, Parrish RK 2nd, Heuer DK, et al. Glaucoma filtering surgery with 5-fluorouracil. *Ophthalmology*. 1987;94(9):1071-8.
36. Franks WA, Hitchings RA. Complications of 5-fluorouracil after trabeculectomy. *Eye (Lond)*. 1991;5 (Pt 4):385-9.
37. (No authors listed). Three-year follow-up of the Fluorouracil Filtering Surgery Study. *Am J Ophthalmol*. 1993;115(1):82-92.
38. Whiteside-Michel J, Liebmann JM, Ritch R. Initial 5-fluorouracil trabeculectomy in young patients. *Ophthalmology*. 1992;99(1):7-13.
39. Wells AP, Cordeiro MF, Bunce C, Khaw PT. Cystic bleb formation and related complications in limbus- versus fornix-based conjunctival flaps in pediatric and young adult trabeculectomy with mitomycin C. *Ophthalmology*. 2003;110(11):2192-7.
40. Okuda T, Higashide T, Fukuhira Y, Kaneko H, Shimomura M, Sugiyama K. Suppression of avascular bleb formation by a thin biodegradable film in a rabbit filtration surgery with mitomycin C. *Graefes Arch Clin Exp Ophthalmol*. 2012;250(10):1441-51.
41. Nilforushan N, Yadgari M, Kish SK, Nassiri N. Subconjunctival bevacizumab versus mitomycin C adjunctive to trabeculectomy. *Am J Ophthalmol*. 2012;153(2):352-7.
42. Nilforushan N, Yadgari M, Falavarjani KG, Afshar AE. Evaluation of subconjunctival oculusgen implantation as an adjunct to trabeculectomy. *Iranian Journal of Ophthalmology*. 2010;22(2):55-62.
43. Nilforushan N, Yadegari M, Hodjat P. Comparison of the success rate of trabeculectomy with OculusGen versus trabeculectomy with mitomycin C. *Iranian Journal of Ophthalmology*. 2011;23(4):3-12.
44. Zahid S, Musch DC, Niziol LM, Lichter PR; Collaborative Initial Glaucoma Treatment Study Group. Risk of endophthalmitis and other long-term complications of trabeculectomy in the Collaborative Initial Glaucoma Treatment Study (CIGTS). *Am J Ophthalmol*. 2013;155(4):674-80.e1.
45. Sony P, Kumar H, Pushker N. Treatment of overhanging blebs with frequency-doubled Nd:YAG laser. *Ophthalmic Surg Lasers Imaging*. 2004;35(5):429-32.
46. Dunnington JH, Regan EF. Late fistulization of operative wounds; diagnosis and treatment. *Trans Am Ophthalmol Soc*. 1949;47:63-79.
47. Wilson MR, Kotas-Neumann R. Free conjunctival patch for repair of persistent late bleb leak. *Am J Ophthalmol*. 1994;117(5):569-74.
48. Sagara H, Sekiryu T, Noji H, Ogasawara M, Imaizumi K, Yago K. Treatment with sodium hyaluronate eye drops in a patient who had early-onset bleb leakage after trabeculectomy with mitomycin C.



- Int Med Case Rep J. 2015;8:301-4.
49. Blok MD, Kok JH, van Mil C, Greve EL, Kijlstra A. Use of the Megasoft Bandage Lens for treatment of complications after trabeculectomy. *Am J Ophthalmol.* 1990;110(3):264-8.
 50. Wu Z, Huang C, Huang Y, Zhang W, Ma D. Soft Bandage Contact Lenses in Management of Early Bleb Leak Following Trabeculectomy. *Eye Sci.* 2015;30(1):13-7.
 51. Kajiwarra K. Repair of a leaking bleb with fibrin glue. *Am J Ophthalmol.* 1990;109(5):599-601.
 52. Leahey AB, Gottsch JD, Stark WJ. Clinical experience with N-butyl cyanoacrylate (Nexacryl) tissue adhesive. *Ophthalmology.* 1993;100(2):173-80.
 53. Spaeth GL. A prospective, controlled study to compare the Scheie procedure with Watson's trabeculectomy. *Ophthalmic Surg.* 1980;11(10):688-94.
 54. Zalta AH, Wieder RH. Closure of leaking filtering blebs with cyanoacrylate tissue adhesive. *Br J Ophthalmol.* 1991;75(3):170-3.
 55. Hill RA, Aminlari A, Sassani JW, Michalski M. Use of a symblepharon ring for treatment of over-filtration and leaking blebs after glaucoma filtration surgery. *Ophthalmic Surg.* 1990;21(10):707-10.
 56. Sugar HS. Complications, repair and reoperation of antiglaucoma filtering blebs. *Am J Ophthalmol.* 1967;63(4):825-33.
 57. Sugar HS. Treatment of hypotony following filtering surgery for glaucoma. *Am J Ophthalmol.* 1971;71(5):1023-33.
 58. Xu JG, Zhong J, Yang YF, Lin MK, Liu X, Yu MB. Efficacy of autologous conjunctival flap on repairing the late-onset filtering bleb leakage. *Int J Ophthalmol.* 2018;11(4):601-6.
 59. Newhouse RP, Beyrer C. Hypotony as a late complication of trabeculectomy. *Ann Ophthalmol.* 1982;14(7):685-6.
 60. Chandler PA. Hypotony after a filtering operation. *Am J Ophthalmol.* 1947;30(4):484.
 61. Baum M, Weiss HS. Argon laser closure of conjunctival bleb leak. *Arch Ophthalmol.* 1993;111(4):438.
 62. Choy BN, Zhu MM, Shum JW, et al. Collagen crosslinking in the management of leaking cystic blebs: a prospective study. *Graefes Arch Clin Exp Ophthalmol.* 2016;254(3):529-33.
 63. Wang Q, Harasymowycz P. Collagen Cross-linking for Late-onset Bleb Leakage: 1-Year Results. *J Glaucoma.* 2016;25(3):e273-e276.
 64. Petursson GJ, Fraunfelder FT. Repair of an inadvertent buttonhole or leaking filtering bleb. *Arch Ophthalmol.* 1979;97(5):926-7.
 65. Tomlinson CP, Belcher CD 3rd, Smith PD, Simmons RJ. Management of leaking filtration blebs. *Ann Ophthalmol.* 1987;19(11):405-11.
 66. Soong HK, Meyer RF, Wolter JR. Fistula excision and peripheral grafts in the treatment of persistent limbal wound leaks. *Ophthalmology.* 1988;95(1):31-6.
 67. Clune MJ, Shin DH, Olivier MM, Kupin TH. Partial-thickness scleral-patch graft in revision of trabeculectomy. *Am J Ophthalmol.* 1993;115(6):818-20.
 68. Kosmin AS, Wishart PK. A full-thickness scleral graft for the surgical management of a late filtration bleb leak. *Ophthalmic Surg Lasers.* 1997;28(6):461-8.
 69. Melamed S, Ashkenazi I, Belcher DC 3rd, Blumenthal M. Donor scleral graft patching for persistent filtration bleb leak. *Ophthalmic Surg.* 1991;22(3):164-5.
 70. Galin MA, Hung PT. Surgical repair of leaking blebs. *Am J Ophthalmol.* 1977;83(3):328-33.
 71. Lee GA, Holcombe DJ. Surgical revision of dysfunctional filtration blebs with bleb preservation, sliding conjunctival flap and fibrin glue. *Eye (Lond).* 2010;24(6):947-53.
 72. Mandal AK. Management of the late leaking filtration blebs. A report of seven cases and a selective review of the literature. *Indian J Ophthalmol.* 2001;49(4):247-54.
 73. Buxton JN, Lavery KT, Liebmann JM, Buxton DE, Ritch R. Reconstruction of filtering blebs with free conjunctival autografts. *Ophthalmology.* 1994;101(4):635-9.
 74. Biswas S, Zaheer I, Monsalve B, Diamond JP.



- Compression sutures with autologous blood injection for leaking trabeculectomy blebs. *Br J Ophthalmol*. 2009;93(4):549-50.
75. Dinah C, Bhachech B, Ghosh G. Long-term success with autologous blood injection for leaking trabeculectomy blebs. *Br J Ophthalmol*. 2010;94(3):392-93.
76. Leen MM, Moster MR, Katz LJ, Terebuh AK, Schmidt CM, Spaeth GL. Management of overfiltering and leaking blebs with autologous blood injection. *Arch Ophthalmol*. 1995;113(8):1050-5.
77. Rauscher FM, Barton K, Budenz DL, Feuer WJ, Tseng SC. Long-term outcomes of amniotic membrane transplantation for repair of leaking glaucoma filtering blebs. *Am J Ophthalmol*. 2007;143(6):1052-4.
78. Maeda H, Eno A, Nakamura M, Negi A. Safe management of a late-onset bleb leak with a needling technique. *Eye (Lond)*. 2000;14(5):802-4.
79. van de Geijn EJ, Lemij HG, de Vries J, de Waard PW. Surgical revision of filtration blebs: a follow-up study. *J Glaucoma*. 2002;11(4):300-5.
80. Busbee BG. Advances in knowledge and treatment: an update on endophthalmitis. *Curr Opin Ophthalmol*. 2004;15(3):232-7.
81. Al-Shahwan S, Al-Torbak AA, Al-Jadaan I, Omran M, Edward DP. Long-term follow up of surgical repair of late bleb leaks after glaucoma filtering surgery. *J Glaucoma*. 2006;15(5):432-6.
82. Lee K, Hyung S. Effect of excision of avascular bleb and advancement of adjacent conjunctiva for treatment of hypotony. *Korean J Ophthalmol*. 2009;23(4):281-5.
83. Budenz DL, Chen PP, Weaver YK. Conjunctival advancement for late-onset filtering bleb leaks: indications and outcomes. *Arch Ophthalmol*. 1999;117(8):1014-9.
84. Budenz DL, Barton K, Tseng SC. Amniotic membrane transplantation for repair of leaking glaucoma filtering blebs. *Am J Ophthalmol*. 2000;130(5):580-8.
85. Hyung SM, Ahn DG. Midterm follow-up of necrotic bleb excision and advancement of the fornical conjunctiva. *Korean J Ophthalmol*. 1999;13(2):85-91.
86. Tannenbaum DP, Hoffman D, Greaney MJ, Caprioli J. Outcomes of bleb excision and conjunctival advancement for leaking or hypotonous eyes after glaucoma filtering surgery. *Br J Ophthalmol*. 2004;88(1):99-103.
87. Wadhvani RA, Bellows AR, Hutchinson BT. Surgical repair of leaking filtering blebs. *Ophthalmology*. 2000;107(9):1681-7.

Footnotes and Financial Disclosures

Conflict of interest:

The authors have no conflict of interest with the subject matter of the present study.